JOURNAL OF NEMATOLOGY

Official Journal of

THE SOCIETY OF NEMATOLOGISTS

VOLUME 11 QUARTERLY IN 1979

Editor-in-Chief
BENJAMIN F. LOWNSBERY

Editorial Board

R. H. Mulvey Joseph A. Veech B. F. Lownsbery R. S. Hussey Howard Ferris

Associate Editors

Donald P. Schmitt Gregory R. Noel
David R. Viglierchio Robert T. Robbins
I. A. Siddiqui Kenneth A. Wright
Kelvin Deming

CONTENTS

JOURNAL OF NEMATOLOGY

Peroxidase Isozymes from Meloidogyne spp. and Their Origin: J. L. Starr Morphological Comparison of Second-Stage Juveniles of Six Populations of Meloidogyne hapla by SEM: J. D. Eisenback and Hedwig Hirschmann Gall Formation on Cirsium arvense by Ditylenchus dipsaci: A. K. Watson and J. D. Shorthouse The Effects of Temperature on Pratylenchus scribneri and P. alleni Populations on Soybeans and Tomatoes: O. J. Dickerson Relative Tollerance of Selected Soybean Cultivars to Hoplolaimus columbus and Possible Effects of Soil Temperature: A. P. Nyczepir and S. A. Lewis Effects of DCPA, EPTC, and Chlorpropham on Pathogenicity of Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra, N. Vovlas, and A. Morgan Golden
Meloidogyne hapla by SEM: J. D. Eisenback and Hedwig Hirschmann Gall Formation on Cirsium arvense by Ditylenchus dipsaci: A. K. Watson and J. D. Shorthouse The Effects of Temperature on Pratylenchus scribneri and P. alleni Populations on Soybeans and Tomatoes: O. J. Dickerson Relative Tollerance of Selected Soybean Cultivars to Hoplolaimus columbus and Possible Effects of Soil Temperature: A. P. Nyczepir and S. A. Lewis Effects of DCPA, EPTC, and Chlorpropham on Pathogenicity of Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
The Effects of Temperature on Pratylenchus scribneri and P. alleni Populations on Soybeans and Tomatoes: O. J. Dickerson Relative Tollerance of Selected Soybean Cultivars to Hoplolaimus columbus and Possible Effects of Soil Temperature: A. P. Nyczepir and S. A. Lewis Effects of DCPA, EPTC, and Chlorpropham on Pathogenicity of Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
Relative Tollerance of Selected Soybean Cultivars to Hoplolaimus columbus and Possible Effects of Soil Temperature: A. P. Nyczepir and S. A. Lewis Effects of DCPA, EPTC, and Chlorpropham on Pathogenicity of Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
and Possible Effects of Soil Temperature: A. P. Nyczepir and S. A. Lewis Effects of DCPA, EPTC, and Chlorpropham on Pathogenicity of Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
Meloidogyne hapla to Alfalfa: G. D. Griffin and J. L. Anderson Control of Meloidogyne javanica and M. arenaria on kenaf and roselle with genetic resistance and nematicides: N. A. Minton and W. C. Adamson Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
Residues of Aldicarb and its Oxides in Beta vulgaris L. and Systemic Control of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
of Heterodera schachtii: Arnold E. Steele Meloidogyne megatyla n. sp., a Root-Knot Nematode from Loblolly Pine: J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
J. G. Baldwin and J. N. Sasser Helicotylenchus oleae n. sp. and H. neopaxilli n. sp. (Hoplolaimidae), Two New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
New Spiral Nematodes Parasitic on Olive Trees in Italy: R. N. Inserra,
The Influence of Temperature on Meloidogyne incognita on Soybean: John F. Nardacci and K. R. Barker
Proleptonchoides southindiae n. gen., n. sp., a New Leptonchoid from South India: V. R. Ferris, C. G. Goseco, and A. C. Kumar
Behavioral Response of <i>Nothanguina phyllobia</i> to Selected Plant Species: A. F. Robinson, C. C. Orr, and J. R. Abernathy
Effect of Oryzalin and 1,1-Dimethylpiperidinium Chloride on Cotton and Tomato Roots Infected with the Root-knot Nematode, <i>Meloidogyne incognita</i> : T. V. Orum, P. G. Bartels, and M. A. McClure
Effects of Management Practices on Nematode and Fungus Populations and Cucumber Yield: A. W. Johnson, D. R. Sumner, and C. A. Jaworski
The Role of Cuticular Strata Nomenclature in the Systematics of Nemata: A. R. Maggenti
RESEARCH NOTES
Techniques for Detecting Dactylella oviparasitica and Evaluating its Significance in Field Soils: G. R. Stirling
Major- and Trace-Element Analyses of a Mermithid Parasite and its Mosquito Host by Proton-induced X-ray Emission: R. Levy, H. A. Van Rinsvelt, and T. W. Miller, Jr.

A Method of Distinguishing Between Living and Dead Nematodes by Enzymatically Induced Fluorescence: Alan F. Bird
pH as a Factor in Parasitism of Mosquito Larvae by the Mermithid Romanomermis culicivorax: James J. Petersen
Equipment and Techniques for Establishing Field Microplots for the Study of Soilborne Pathogens: K. R. Barker, B. I. Daughtry, and D. W. Corbett
New Data for Leptonchus transvaalensis from Nigeria, and Key to Leptonchus: C. G. Goseco and V. R. Ferris
Immune Responses of Mosquitoes against Romanomermis culicivorax (Mermithida: Nematoda): George O. Poinar, Jr., Roberta T. Hess, and James J. Petersen
Vol. 11, No. 2 April 19
The Histopathological Reactions of Vigna sinensis to Separate and Concomitant Parasitism by Meloidogyne javanica and Rotylenchulus reniformis: A. H. Y. Taha and A. S. Kassab
The Relationship Between Population Density of Heterodera schachtii, Soil Temperature, and Sugarbeet Yields: D. A. Cooke and I. J. Thomason
Pathogenicity of Hirschmanniella oryzae, H. spinicaudata, and H. imamuri on Rice: J. O. Babatola and John Bridge
Interaction of Population Levels of Fusarium oxysporum f. sp. vasinfectum and Meloidogyne incognita on Cotton: R. H. Garber, E. C. Jorgenson, S. Smith, and A. H. Hyer
Chemical Control of Nematodes and Soil-borne Plant-Pathogenic Fungi on Cabbage Tranplants: A. W. Johnson, S. M. McCarter, C. A. Jaworski, and R. E. Williamson
Granular Nematicides as Adjuncts to Fumigants for Control of Cotton Root-knot Nematodes: E. C. Jorgenson
Effects of Nematicide Placement on Nematode Populations and Soybean Yields: N. A. Minton, M. B. Parker, O. L. Brooks, and C. E. Perry
Effects of Meloidogyne incognita on Nitrogen Fixation in Soybean: J. G. Baldwin, K. R. Barker, and L. A. Nelson
Growth Reduction of Apple Seedlings by <i>Pratylenchus penetrans</i> as Influenced by Seedling Age at Inoculation: B. A. Jaffee and W. F. Mai
Effect of Soil Water Potential on Growth of Apple Trees Infected with Pratylenchus penetrans: B. A. Jaffee and W. F. Mai
Quantitative Aspects of the Development of <i>Meloidogyne arenaria</i> Larvae in Grapevine Varieties and Rootstocks: H. Ferris and W. A. Hunt
Descriptions of <i>Meloidogyne camelliae</i> n.sp. and <i>M. querciana</i> n.sp (Nematoda:Meloidogynidae), with SEM and Host-Range Observations: A. Morgan Golden
Anhydrobiotic Coiling of Nematodes in Soil: Y. Demeure, D. W. Freckman, and S. D. Van Gundy

iv Journal of Nematology, Volume 11, No. 4, October 1979	
RESEARCH NOTES	
Modification of the Mini-sieve and Prepolymerized Plate Technique in Electron Microscopy: A. De Grisse	
Influence of Plant Age, Light Intensity, Nematode Inoculun and Their Interactions on Tomato Growth and Reprodu <i>Meloidogyne hapla:</i> B. Fawole and W. F. Mai	ction o
Technique for Screening Cowpea Germplasm for Resistance to R Nematodes: J. R. Rich, H. E. McKinney, and I. J. Thomason	
A Rapid Method for Sampling Surface Soil: D. A. Cooke, H. E. M and I. J. Thomason	
Chemical Control of <i>Heterodera schachtii</i> on Sugarbeet in Californ Cooke, I. J. Thomason, H. E. McKinney, W. E. Bendixen, an Hagemann	d R. W
Automated Dehydration of Small Specimens: L. J. Stowell and Mi McClure	
Nonsolanaceous Hosts of <i>Globodera</i> in the Andes: Parviz Jatala, J. A. Vilca, and W. Cornejo	
/ol 11 No 2	Inly '
Vol. 11, No. 3	-
Vol. 11, No. 3 Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek	pecies of
Soil Nematodes in Terrestrial Ecosystems: G. W. YeatesInfluence of Temperature on Population Development of Eight S	pecies of
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant	pecies of and Re- Croll
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant pellent and in Thermal Gradients: T. A. Rutherford and N. A. Histochemical Localization and Nematoxicity of Terpenoid Alde	and Re- Croll
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant pellent and in Thermal Gradients: T. A. Rutherford and N. A. Histochemical Localization and Nematoxicity of Terpenoid Alde Cotton: J. A. Veech The Influence of Glomus mosseae on Tylenchulus semipenetrans— and Uninfected Citrus limon Seedlings: J. H. O'Bannon, R. N.	and Re- Croll hydes in Infected Inserra
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant pellent and in Thermal Gradients: T. A. Rutherford and N. A. Histochemical Localization and Nematoxicity of Terpenoid Alde Cotton: J. A. Veech The Influence of Glomus mosseae on Tylenchulus semipenetrans— and Uninfected Citrus limon Seedlings: J. H. O'Bannon, R. N. S. Nemec and N. Vovlas Developmental Nutrition of Nematodes: the Biochemical Role of Heme Compounds, and Lysosomal Enzymes: R. Bolla	and Re- Croll
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant pellent and in Thermal Gradients: T. A. Rutherford and N. A. Histochemical Localization and Nematoxicity of Terpenoid Alde Cotton: J. A. Veech The Influence of Glomus mosseae on Tylenchulus semipenetrans— and Uninfected Citrus limon Seedlings: J. H. O'Bannon, R. N. S. Nemec and N. Vovlas Developmental Nutrition of Nematodes: the Biochemical Role of Heme Compounds, and Lysosomal Enzymes: R. Bolla Some Factors Causing Variability in 1,2-Dibromo-3-Chloropropane	and Re- Croll hydes in Infected Inserra Sterols,
Soil Nematodes in Terrestrial Ecosystems: G. W. Yeates Influence of Temperature on Population Development of Eight S Pratylenchus on Soybean: Nelia Acosta and R. B. Malek Wave Forms of Caenorhabditis elegans in a Chemical Attractant pellent and in Thermal Gradients: T. A. Rutherford and N. A. Histochemical Localization and Nematoxicity of Terpenoid Alde Cotton: J. A. Veech The Influence of Glomus mosseae on Tylenchulus semipenetrans— and Uninfected Citrus limon Seedlings: J. H. O'Bannon, R. N. S. Nemec and N. Vovlas Developmental Nutrition of Nematodes: the Biochemical Role of Heme Compounds, and Lysosomal Enzymes: R. Bolla Some Factors Causing Variability in 1,2-Dibromo-3-Chloropropane trations in Soil: M. V. McKenry and P. Naylor Effects of Phenamiphos, Methyl Bromide, and Fallowing on Prat penetrans, Yield of Medicago sativa, and Fusarium Infection	and Re-Croll hydes in Infected Inserra, Concentration of the Concentrati

Mode of Parasitism of *Meloidogyne* and Other Nematode Eggs by *Dactylella oviparasitica*: G. R. Stirling and R. Mankau

282

RESEARCH NOTE

Interacting Effects of Soil Temperature and Type on Reproduction and Pathogenicity of <i>Heterodera schachtii</i> and <i>Meloidogyne hapla</i> on Sugarbeets: G. S. Santo and W. J. Bolander	289
Vol. 11, No. 4 October 1	979
Abstracts of Papers Presented at the Eighteenth Annual Meeting of the Society of Nematologists, Salt Lake City, Utah, July 23-26, 1979	293
Pathogenicity of Macrophomina phaseoli on Jute in the Presence of Meloidogyne incognita and Hoplolaimus indicus: MD. Samsul Haque and M. C. Mukhopadhyaya	318
Description and Bionomics of Mesomermis camdenensis n. sp. (Mermithidae), a Parasite of Black Flies (Simuliidae): Daniel Molloy	321
Nonfumigant Nematicides for Control of Root-knot Nematode to Protect Carrot Root Growth in Organic Soils: T. C. Vrain, G. Belair, and P. Martel	328
Helicotylenchus stylocercus n. sp. and Rotylenchus phaliurus n. sp. (Nematoda: Hoplolaimidae) from Costa Rica: M. R. Siddiqi and J. Pinochet	333
Histological Study of the Compatible and Incompatible Interaction of Soybeans and <i>Meloidogyne incognita</i> : D. T. Kaplan, I. J. Thomason, and S. D. Van Gundy	338
New Species of Bunonematoidea (Rhabditida) from Georgia and Tennessee: E. C. Bernard	344
Partial Characterization of the Cuticle Surface of <i>Meloidogyne javanica</i> Females: S. Himmelhoch, D. Orion, and B. M. Zuckerman	358
Heterodera canadensis n. sp. (Nematoda: Heteroderidae) from Spike-Rush (Eleocharis acicularis (L.) R. & S.) in Quebec, Canada: Roland H. Mulvey	363
Polyploidy in an Amphimictic Population of Heterodera glycines: A. C. Triantaphyllou and R. D. Riggs	371
Response of <i>Pinus ponderosa</i> Seedlings to Stylet-Bearing Nematodes: D. R. Viglierchio	377
Control of Tylenchulus semipenetrans on Citrus With Aldicarb, Oxamyl, and DBCP: L. W. Timmer and J. V. French	387
	395
RESEARCH NOTE	•
Vegetable Oils as Protectants Against Nematode Infections: P. M. Miller	402
INDEX FOR VOLUME 11 (1979)	2
TABLE OF CONTENTS FOR VOLUME 11 (1979)	i-v

